

Multiple Solutions for a Semilinear Elliptic Equation

Author(s): Manuel A. Del Pino and Patricio L. Felmer

Source: *Transactions of the American Mathematical Society*, Vol. 347, No. 12 (Dec., 1995), pp. 4839-4853

Published by: [American Mathematical Society](#)

Stable URL: <http://www.jstor.org/stable/2155065>

Accessed: 12-07-2015 00:05 UTC

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



American Mathematical Society is collaborating with JSTOR to digitize, preserve and extend access to *Transactions of the American Mathematical Society*.

<http://www.jstor.org>

MULTIPLE SOLUTIONS FOR A SEMILINEAR ELLIPTIC EQUATION

MANUEL A. DEL PINO AND PATRICIO L. FELMER

ABSTRACT. Let Ω be a bounded, smooth domain in \mathbb{R}^N , $N \geq 1$. We consider the problem of finding nontrivial solutions to the elliptic boundary value problem

$$\begin{aligned}\Delta u + \lambda u &= h(x)|u|^{p-1}u \quad \text{in } \Omega, \\ u &= 0 \quad \text{on } \partial\Omega,\end{aligned}$$

where $h \geq 0$, $h \not\equiv 0$ is Hölder continuous on $\bar{\Omega}$ and $p > 1$, λ are constants.

Let Ω_0 denote the interior of the set where h vanishes in Ω . We assume $h > 0$ a.e. on $\Omega \setminus \Omega_0$ and consider the eigenvalues $\lambda_i(\Omega)$ and $\lambda_i(\Omega_0)$ of the Dirichlet problem in Ω and Ω_0 respectively. We prove that no nontrivial solution of the equation exists if λ satisfies, for some $k \geq 1$,

$$\lambda_k(\Omega_0) \leq \lambda \leq \lambda_{k+1}(\Omega).$$

On the other hand, if, for some nonnegative integers s, k with $s \geq k + 1$, λ satisfies

$$\lambda_s(\Omega) < \lambda < \lambda_{k+1}(\Omega_0),$$

then the equation above possesses at least $s - k$ pairs of nontrivial solutions.

For the proof of these results we use a variational approach. In particular, the existence result takes advantage of the even character of the associated functional.

0. INTRODUCTION

Let Ω be a bounded, smooth domain in \mathbb{R}^N , $N \geq 1$. We consider the problem of finding nontrivial solutions to the elliptic boundary value problem

$$(0.1) \quad \Delta u + \lambda u = h(x)|u|^{p-1}u \quad \text{in } \Omega,$$

$$(0.2) \quad u = 0 \quad \text{on } \partial\Omega,$$

where h is a nonnegative, not identically zero Hölder continuous function defined on $\bar{\Omega}$ and $p > 1$, λ are constants.

Ouyang [5] studied existence and uniqueness of positive solutions to problem (0.1)–(0.2) and also of its corresponding analogue on a compact Riemannian manifold. That paper mainly focused on studying the effect of *vanishing* of the

Received by the editors October 13, 1993 and, in revised form, October 24, 1994.

1991 *Mathematics Subject Classification.* Primary 35J25, 35J20.

Key words and phrases. Variational method, multiplicity, uniqueness, blow up of solutions.

The work of the first author was partially supported by NSF grant DMS-9100383.

The work of the second author has been partially supported by FONDECYT under grant No 1212-91.

function h , a question originally raised by Kazdan and Warner [4] who considered the problem on a compact manifold and proved, among other results, that for strictly positive h there is a unique positive solution for any $\lambda > 0$. Similarly, a unique positive solution to (0.1)–(0.2) exists for any $\lambda > \lambda_1(\Omega)$, where $\lambda_1(\Omega)$ denotes the first Dirichlet eigenvalue of the Laplacian in Ω . An integration by parts easily yields that this condition is also necessary for existence of a nontrivial solution to (0.1)–(0.2).

Let Ω_0 denote the interior of the set where h vanishes in Ω . We will assume henceforth that Ω_0 is nonempty and that $h > 0$ a.e. on $\Omega \setminus \Omega_0$. Then the following result holds; see [5] and also [3] for an alternative proof.

Theorem 0.1. *There exists a unique positive solution u_λ to problem (0.1)–(0.2) if $\lambda_1(\Omega) < \lambda < \lambda_1(\Omega_0)$. Moreover,*

$$\lim_{\lambda \rightarrow \lambda_1(\Omega_0)} \|u_\lambda\|_{L^2} = +\infty$$

and no positive solution exists in case that $\lambda \geq \lambda_1(\Omega_0)$.

Ouyang actually proves that the set of all pairs (λ, u) with u a positive solution of (0.1)–(0.2) is constituted exactly by the first bifurcation branch starting at $(\lambda_1(\Omega), 0)$ and that this branch continues to exist until λ reaches $\lambda_1(\Omega_0)$ where it blows up. We should remark that a very different behavior of this branch takes place in case that h changes sign (see [6]).

Now, it follows from standard bifurcation theory that a branch of nontrivial solutions of (0.1)–(0.2) starts at each eigenvalue of the Laplacian, provided that this eigenvalue is e.g. simple. Thus, the natural question arises: Will branches associated with higher eigenvalues exhibit a similar behavior to that of the first one? This paper is mainly motivated by this question. Giving a complete answer may be very hard. However, our first result, Theorem 0.2 below, partially recovers the picture of the first branch: it shows, in particular, that if the Dirichlet eigenvalues of both Ω and Ω_0 are simple, then for each positive integer k there exists a family of pairs $(\lambda, \pm u_\lambda^k)$ with u_λ^k a nontrivial solution of (0.1)–(0.2), defined for $\lambda \in (\lambda_k(\Omega), \lambda_k(\Omega_0))$, with the property that $\|u_\lambda^k\|_{L^2}$ blows up as λ approaches $\lambda_k(\Omega_0)$. Moreover, these “branches” do not intersect each other in the sense that, if λ belongs to j intervals of the form $(\lambda_k(\Omega), \lambda_k(\Omega_0))$ then at least j pairs of nontrivial solutions exist.

To state our first result in the general case, we consider the sequences of Dirichlet eigenvalues for Ω and Ω_0 ,

$$\begin{aligned} \lambda_1(\Omega) < \lambda_2(\Omega) \leq \lambda_3(\Omega) \leq \lambda_4(\Omega) \leq \dots, \\ \lambda_1(\Omega_0) < \lambda_2(\Omega_0) \leq \lambda_3(\Omega_0) \leq \dots \end{aligned}$$

where a given eigenvalue in the above sequences appears as many times as its multiplicity indicates. If Ω_0 is a general open subset of Ω , with no regularity assumed on its boundary, its Dirichlet eigenvalues can be naturally defined considering the closed subspace $H_*^1(\Omega_0)$ of $H_0^1(\Omega)$ given by

$$H_*^1(\Omega_0) = \{u \in H_0^1(\Omega) \mid u = 0 \text{ a.e. on } \Omega \setminus \Omega_0\}.$$

It is well known that $H_*^1(\Omega_0) = H_0^1(\Omega_0)$ if Ω_0 has, for example, a Lipschitz continuous boundary. We define the sequence of Dirichlet eigenvalues of Ω_0 as those of the problem

$$u = \lambda T u, \quad u \in H_*^1(\Omega_0),$$

where T is the compact, selfadjoint operator so that $v = Tf$ is the unique element of $H_*^1(\Omega_0)$ satisfying

$$\int_{\Omega_0} \nabla v \nabla \phi = \int_{\Omega_0} f \phi \quad \forall \phi \in H_*^1(\Omega_0).$$

Note that the variational characterization of the eigenvalues yields

$$(0.3) \quad \lambda_k(\Omega) \leq \lambda_k(\Omega_0) \quad \forall k \geq 1.$$

Our first result is the following

Theorem 0.2. *Assume that there exist nonnegative integers s, k with $s \geq k + 1$ such that λ satisfies*

$$(0.4) \quad \lambda_s(\Omega) < \lambda < \lambda_{k+1}(\Omega_0).$$

Then there exist at least $(s - k)$ distinct pairs of nontrivial solutions $\pm u_\lambda^{k+1}, \pm u_\lambda^{k+2}, \dots, \pm u_\lambda^s$ to (0.1)–(0.2). Moreover, if q is the multiplicity of $\lambda_{k+1}(\Omega_0)$, then the above families can be selected in such a way that

$$(i) \quad \lim_{\lambda \rightarrow \lambda_{k+1}(\Omega_0)} \|u_\lambda^l\|_{L^2} = +\infty, \quad k + 1 \leq l \leq \min\{s, k + q\},$$

and

$$(ii) \quad \|u_\lambda^l\|_{L^2} \text{ remains uniformly bounded on } \lambda \text{ satisfying (0.4) for all } l \text{ satisfying } k + q < l \leq s, \text{ in case that } q < s - k.$$

Observe that among the assumptions of the above result we may also include, without loss of generality,

$$\lambda_{s+1}(\Omega) > \lambda_s(\Omega) \quad \text{and} \quad \lambda_k(\Omega_0) < \lambda_{k+1}(\Omega_0),$$

in which case the number of families predicted by the theorem becomes maximal.

Let us notice that if the eigenvalues of Ω_0 are simple, this result provides, for each $k \geq 1$, a family of pairs of nontrivial solutions u_λ^l , defined for $\lambda_k(\Omega) < \lambda < \lambda_{k+1}(\Omega_0)$, which blows up in L^2 -norm as λ approaches $\lambda_{k+1}(\Omega_0)$.

It is natural to ask what happens if λ does not belong to any interval of the real line covered by the above theorem. It is maybe a little surprising that a full answer is possible in this situation: no nontrivial solution of (0.1)–(0.2) exists.

Theorem 0.3. *Assume that λ satisfies, for some $k \geq 1$,*

$$\lambda_k(\Omega_0) \leq \lambda \leq \lambda_{k+1}(\Omega).$$

Then (0.1)–(0.2) possesses only the solution $u \equiv 0$.

The proofs of Theorems 0.2 and 0.3 are based on variational arguments applied to the functional

$$(0.5) \quad J_\lambda(u) = \frac{1}{2} \int_{\Omega} |\nabla u|^2 - \lambda u^2 + \frac{1}{p+1} \int_{\Omega} h|u|^{p+1}$$

defined on the Banach space

$$(0.6) \quad X = \left\{ u \in H_0^1(\Omega) \mid \int_{\Omega} h|u|^{p+1} < +\infty \right\}$$

endowed with norm

$$(0.7) \quad \|u\|_X = \left(\int_{\Omega} |\nabla u|^2 \right)^{1/2} + \left(\int_{\Omega} h|u|^{p+1} \right)^{\frac{1}{p+1}}.$$

It is standard to check that J_{λ} is of class C^1 on X and that its critical points correspond exactly to the solutions of (0.1)–(0.2). We should remark that the positive solutions correspond precisely to global minimizers of this functional as established in [3]. If $\lambda \geq \lambda_1(\Omega_0)$ the functional becomes unbounded below. We will take advantage of the even character of the functional J_{λ} in order to establish existence of multiple critical points under the assumptions of Theorem 0.2. Properties (i) and (ii) will follow from appropriate min-max characterizations of the associated critical values. Actually, these characterizations give that $J_{\lambda}(u'_{\lambda}) \rightarrow -\infty$ as λ approaches $\lambda_{k+1}(\Omega_0)$, in case (i) and remains locally uniformly bounded for λ in the range given by (ii).

On the other hand, the result of Theorem 0.3 actually holds true for a much larger class of nonlinearities. In particular, no symmetry is required. See Remark 1.1 below.

The rest of this paper will be devoted to the proof of the above results. Its outline is as follows. In §1 we prove the nonexistence result of Theorem 0.3. In §2 we show that J_{λ} satisfies the Palais-Smale condition for any value of λ , a necessary step for the proof of Theorem 0.2, which we carry out in §3.

In what remains of this paper X will always denote the space given by (0.6) with its associated norm. In the space $H_0^1(\Omega)$ we consider the usual inner product

$$\langle \psi, \phi \rangle = \int_{\Omega} \nabla \psi \nabla \phi, \quad \psi, \phi \in H_0^1(\Omega),$$

and denote by $\|\cdot\|_{H_0^1}$ its associated norm. Correspondingly, we denote by $\|\cdot\|_{L^2}$ the usual norm in $L^2(\Omega)$.

We also choose orthogonal bases of $H_0^1(\Omega)$ and $H_*^1(\Omega_0)$ respectively denoted by $\{\psi_i\}_{i=1}^{\infty}$ and $\{\phi_i\}_{i=1}^{\infty}$ such that ψ_i is an eigenfunction associated to $\lambda_i(\Omega)$ and ϕ_i an eigenfunction for $\lambda_i(\Omega_0)$, normalized so that $\|\phi_i\|_{L^2} = \|\psi_i\|_{L^2} = 1$.

1. PROOF OF THEOREM 0.3

In this section we shall prove that if for some $k \geq 1$ one has

$$(1.1) \quad \lambda_k(\Omega_0) \leq \lambda \leq \lambda_{k+1}(\Omega),$$

then no nontrivial solution of (0.1)–(0.2) exists. We begin with a preliminary result.

Lemma 1.1. *For any $\varepsilon > 0$ there exist functions:*

(a) $\psi_1^{\varepsilon}, \psi_2^{\varepsilon}, \dots, \psi_k^{\varepsilon} \in H_0^1(\Omega)$ such that $\|\psi_i - \psi_i^{\varepsilon}\|_{H_0^1} < \varepsilon$ for $i = 1, \dots, k$ and

$$(1.2) \quad \text{span}\{\phi_1, \dots, \phi_k\} \cap \text{span}\{\psi_1^{\varepsilon}, \dots, \psi_k^{\varepsilon}\}^{\perp} = \{0\}, \text{ and}$$

(b) $\phi_1^{\varepsilon}, \phi_2^{\varepsilon}, \dots, \phi_k^{\varepsilon} \in H_*^1(\Omega_0)$ such that $\|\phi_i - \phi_i^{\varepsilon}\|_{H_0^1} < \varepsilon$ for $i = 1, \dots, k$ and

$$(1.3) \quad \text{span}\{\phi_1^{\varepsilon}, \dots, \phi_k^{\varepsilon}\} \cap \text{span}\{\psi_1, \dots, \psi_k\}^{\perp} = \{0\}.$$

Here, \perp denotes the orthogonal in $H_0^1(\Omega)$.

Proof. First we prove (a). We need to choose functions $\eta_1^\varepsilon, \dots, \eta_k^\varepsilon$ with $\|\eta_i^\varepsilon\|_{H_0^1} < \varepsilon$ such that if we set $\psi_i^\varepsilon = \psi_i + \eta_i^\varepsilon$, then the matrix

$$A_\varepsilon = \{\langle \psi_i^\varepsilon, \phi_j \rangle\}_{i,j}$$

is invertible. Observe that we can find numbers λ_{ij} with $\sum_{i,j} \lambda_{ij}^2$ arbitrarily small, so that the matrix with coefficients $\langle \psi_i, \phi_j \rangle + \lambda_{ij}$ is invertible. Let us set $\eta_i^\varepsilon = \sum_{j=1}^k \lambda_{ij} \phi_j$. For appropriately chosen λ_{ij} , the η_i^ε defined in this way will clearly satisfy the desired properties.

For the proof of part (b), we need to find functions $\eta_1^\varepsilon, \dots, \eta_k^\varepsilon \in H_*^1(\Omega_0)$ with $\|\eta_i^\varepsilon\|_{H_0^1} < \varepsilon$ such that the matrix with coefficients $\langle \psi_i, \phi_j + \eta_j^\varepsilon \rangle$ is invertible. Note first that the matrix with coefficients $\int_{\Omega_0} \psi_i \psi_j$ is invertible. Indeed, this is a consequence of the fact that the functions ψ_1, \dots, ψ_k are linearly independent on Ω_0 , as follows from the unique continuation property for a linear combination of them. Now choose functions $\tilde{\psi}_1, \dots, \tilde{\psi}_k \in C_0^\infty(\Omega_0)$ such that the matrix with coefficients $\int_{\Omega_0} \psi_i \tilde{\psi}_j$ is still invertible. Finally, as in part (a), η_i^ε defined as some linear combination of the $\tilde{\psi}_j$'s with sufficiently small coefficients will satisfy the desired requirement. \square

Proof of Theorem 0.3. First we consider the case

$$\lambda_k(\Omega_0) < \lambda < \lambda_{k+1}(\Omega).$$

For a small number ε to be chosen, we consider functions $\psi_1^\varepsilon, \dots, \psi_k^\varepsilon$ as in Lemma 1.1, part (a). Let us set $V = \text{span}\{\phi_1, \dots, \phi_k\}$ and

$$Y = \{y \in X \mid \langle y, \psi_i^\varepsilon \rangle = 0, i = 1, \dots, k\}.$$

For $v = \sum_{i=1}^k t_i \phi_i$ and $y \in Y$ we see that

$$J_\lambda(v + y) = J_\lambda(y) + \sum_{i=1}^k \frac{t_i^2}{2} (\lambda_i(\Omega_0) - \lambda) + \sum_{i=1}^k t_i \int_{\Omega} (\nabla \phi_i \nabla y - \lambda \phi_i y).$$

Let $\tilde{u} = \tilde{v} + \tilde{y}$ be a critical point of J_λ . We will show that $\tilde{u} \equiv 0$. To do this, we see that, for each fixed $y \in Y$, the functional $v \mapsto J_\lambda(v + y)$ has a unique critical point $v = w(y) = \sum_{i=1}^k t_i(y) \phi_i$ where

$$(1.4) \quad t_i(y) = - \frac{\int_{\Omega} \nabla \phi_i \nabla y - \lambda \phi_i y}{\lambda_i(\Omega_0) - \lambda}, \quad i = 1, \dots, k.$$

Thus, we must have $\tilde{v} = w(\tilde{y})$. Moreover, \tilde{y} must be a critical point of the functional

$$(1.5) \quad y \mapsto J_\lambda(w(y) + y) = J_\lambda(y) + \frac{1}{2} \sum_{i=1}^k \frac{(\int_{\Omega} \nabla \phi_i \nabla y - \lambda \phi_i y)^2}{\lambda - \lambda_i(\Omega_0)}.$$

We will check that the above functional is strictly convex if ε is chosen sufficiently small. To do this, it clearly suffices to show that the quadratic functional

$$(1.6) \quad I_\lambda(y) = \int_{\Omega} |\nabla y|^2 - \lambda \int_{\Omega} |y|^2$$

is strictly convex on Y , which is equivalent to verifying that $I_\lambda(y) > 0$ for $y \neq 0$. Let $\bar{y} = \sum_{i=1}^k \langle y, \psi_i \rangle \psi_i$, $x = y - \bar{y}$. Then $I_\lambda(y) = I_\lambda(\bar{y}) + I_\lambda(x)$, and since $\lambda < \lambda_{k+1}(\Omega)$, we have

$$(1.7) \quad I_\lambda(y) \geq \left(1 - \frac{\lambda}{\lambda_{k+1}(\Omega)}\right) \|x\|_{H_0^1}^2 + \left(1 - \frac{\lambda}{\lambda_1(\Omega)}\right) \|\bar{y}\|_{H_0^1}^2.$$

But

$$(1.8) \quad \|\bar{y}\|_{H_0^1}^2 = \sum_{i=1}^k |\langle y, \psi_i \rangle|^2 = \sum_{i=1}^k |\langle y, \psi_i - \psi_i^\varepsilon \rangle|^2 \leq k\varepsilon^2 \|y\|_{H_0^1}^2$$

and hence

$$(1.9) \quad \|x\|_{H_0^1}^2 = \|y\|_{H_0^1}^2 - \|\bar{y}\|_{H_0^1}^2 \geq (1 - k\varepsilon^2) \|y\|_{H_0^1}^2.$$

Combining (1.7), (1.8) and (1.9) we obtain, for ε sufficiently small,

$$I_\lambda(y) > 0 \quad \text{if } y \neq 0.$$

It follows that the functional (1.5) has $y = 0$ as its unique critical point, so that $\tilde{x} = 0$ and hence $u = 0$, as desired.

Next let us assume that $\lambda = \lambda_k(\Omega_0)$ and let q be the multiplicity of $\lambda_k(\Omega_0)$. In this case, for each $y \in Y$, the functional $v \mapsto J_\lambda(y + v)$ has a critical point if and only if

$$\int_\Omega \nabla \phi_l \nabla y - \lambda \int_\Omega \phi_l y = 0, \quad k \leq l \leq k - q + 1.$$

And in such case the critical points are given by the q -parameter family

$$v_s(y) = \sum_{i=1}^{k-q} t_i(y) \phi_i + \sum_{i=k-q+1}^k s_i \phi_i, \quad s = (s_{k-q+1}, \dots, s_k) \in \mathbb{R}^q,$$

where $t_i(y)$ is as in (1.4). Then if $\tilde{u} = \tilde{v} + \tilde{y}$ is a critical point of J_λ , then $\tilde{v} = v_{\tilde{s}}(\tilde{y})$ for some $\tilde{s} \in \mathbb{R}^q$, and \tilde{y} must be a critical point of the functional $y \mapsto J_\lambda(y + v_{\tilde{s}}(y))$. But, similarly as in the previous case, an appropriate choice of ε yields that $y = 0$ is the only critical point of this functional. Thus, $\tilde{u} = \sum_{i=k-q+1}^k \tilde{s}_i \phi_i$ is a critical point of J_λ and hence satisfies

$$(1.10) \quad \Delta \tilde{u} + \lambda_k(\Omega_0) \tilde{u} = 0 \quad \text{in } \Omega.$$

The unique continuation property of the solutions of (1.10) yields that $\tilde{u} = 0$, as desired.

It only remains to consider the case $\lambda = \lambda_{k+1}(\Omega)$. Let $\phi_1^\varepsilon, \dots, \phi_k^\varepsilon$ be as in Lemma 1.1, part (b) and consider the decomposition $X = V \oplus Y$ where now $V = \text{span}\{\phi_1^\varepsilon, \dots, \phi_k^\varepsilon\}$, and

$$Y = \{y \in X \mid \langle y, \psi_i \rangle = 0, i = 1, \dots, k\}.$$

For $v = \sum_{i=1}^k t_i \phi_i^\varepsilon$ and $y \in Y$ we now have that

$$(1.11) \quad J_\lambda(v + y) = J_\lambda(y) + \sum_{i=1}^k \frac{t_i^2}{2} (\lambda_i(\Omega_0) - \lambda) + \sum_{i=1}^k t_i \left(\int_\Omega \nabla \phi_i^\varepsilon \nabla y - \lambda \phi_i^\varepsilon y \right)$$

$$(1.12) \quad + \frac{1}{2} \sum_{i=1}^k \sum_{j=1}^k t_i t_j o_{ij}(\varepsilon)$$

where

$$o_{ij}(\varepsilon) = \int_\Omega \nabla(\phi_i + \eta_i^\varepsilon) \nabla \eta_j^\varepsilon - \lambda(\phi_i + \eta_i^\varepsilon) \eta_j^\varepsilon$$

with $\eta_i^\varepsilon = \phi_i - \phi_i^\varepsilon$. From the choice of the functions ϕ_i^ε in Lemma 1.1, part (b) we certainly have that $\lim_{\varepsilon \rightarrow 0} o_{ij}(\varepsilon) = 0$ for all $1 \leq i, j \leq k$. Recalling that, for $1 \leq i \leq k$, $\lambda_i(\Omega_0) - \lambda < 0$ we see that for $\varepsilon > 0$ sufficiently small, as in the previous cases, $\tilde{u} = \tilde{v} + \tilde{y}$ will be a critical point of J_λ if and only if $\tilde{v} = w(\tilde{y}) = \sum_{i=1}^k t_i(\tilde{y}) \phi_i^\varepsilon$ with $t(y) = (t_1(y), \dots, t_k(y))$ given by

$$t(y) = -(D + O(\varepsilon))^{-1} b(y)$$

where $D = \text{diag}(\lambda_i(\Omega_0) - \lambda)$, $O(\varepsilon) = (o_{ij}(\varepsilon))_{ij}$ and $b(y) = (b_1(y), \dots, b_k(y))$ is the vector with components

$$b_i(y) = \int_\Omega \nabla \phi_i^\varepsilon \nabla y - \lambda \phi_i^\varepsilon y.$$

Moreover, \tilde{y} is a critical point of the functional

$$(1.13) \quad y \mapsto J_\lambda(w(y) + y) = J_\lambda(y) - \frac{1}{2} b(y)^\tau (D + O(\varepsilon))^{-1} b(y).$$

But this functional is convex since $\int_\Omega |\nabla y|^2 - \lambda \int_\Omega |y|^2 \geq 0$ and $D + O(\varepsilon)$ is negative definite for ε small enough. Thus, its critical points are all minima and then for \tilde{y} , a minimum of (1.13), we have

$$\int_\Omega |\nabla \tilde{y}|^2 - \lambda \int_\Omega |\tilde{y}|^2 = 0.$$

Hence $\tilde{y} = \sum_{i=1}^q \tilde{t}_i \psi_{k+i}$ for some $\tilde{t} = (t_1, \dots, t_q) \in \mathbb{R}^q$, where q is the multiplicity of $\lambda_{k+1}(\Omega)$. Also, at the minimum \tilde{y} we have

$$\frac{1}{p+1} \int_\Omega h |\tilde{y}|^{p+1} = 0;$$

hence \tilde{y} has its support contained in Ω_0 . The unique continuation property of the eigenfunctions $\{\psi_i\}$ thus yields $\tilde{y} = 0$, and therefore $\tilde{u} \equiv 0$ as desired. This concludes the proof. \square

Remark 1.1. The preceding arguments actually apply to the more general problem

$$(1.14) \quad \Delta u + \lambda u = h(x) f(u) \quad \text{in } \Omega,$$

$$(1.15) \quad u = 0 \quad \text{on } \partial\Omega,$$

where we assume that f is locally Hölder continuous, nondecreasing and such that $uf(u) > 0$ for $u \neq 0$. In fact, exactly the same proof yields that if

$$\lambda_k(\Omega_0) \leq \lambda \leq \lambda_{k+1}(\Omega),$$

then (1.14)–(1.15) possesses only the solution $u \equiv 0$.

2. THE PALAIS-SMALE CONDITION

The proof of Theorem 0.2 involves variational arguments applied to the functional J_λ defined on the space X given by (0.6) as

$$J_\lambda(u) = \frac{1}{2} \int_\Omega |\nabla u|^2 - \lambda u^2 + \frac{1}{p+1} \int_\Omega h|u|^{p+1}.$$

In this section we will show that the Palais-Smale condition holds for J_λ for any value of λ .

Lemma 2.1. *J_λ satisfies the Palais-Smale condition for any λ ; that is, any sequence $\{u_n\} \subset X$ such that $J_\lambda(u_n)$ is bounded and $J'_\lambda(u_n) \rightarrow 0$ in X^* possesses a convergent subsequence in X .*

Proof. As a first step we show that the sequence $\{u_n\}$ as in the statement of the lemma is bounded in X . Assume $J_\lambda(u_n) \rightarrow C \in \mathbb{R}$ or

$$(2.1) \quad \frac{1}{2} \int_\Omega |\nabla u_n|^2 - \lambda u_n^2 + \frac{1}{p+1} \int_\Omega h|u_n|^{p+1} = C + o(1).$$

We also have

$$(2.2) \quad \int_\Omega \nabla u_n \nabla \phi - \lambda \int_\Omega u_n \phi + \int_\Omega h|u_n|^{p-1} u_n \phi = o(1) \|\phi\|_X$$

for all $\phi \in X$ where $o(1) \rightarrow 0$ uniformly on ϕ .

Let us assume, by contradiction, $\|u_n\|_X \rightarrow +\infty$. Then, from (2.1) we see that, for some $c > 0$,

$$(2.3) \quad \|u_n\|_X \leq c \|u_n\|_{L^2}$$

for large n .

Let us set $\hat{u}_n = u_n / \|u_n\|_{L^2}$. Then, from (2.3), \hat{u}_n is bounded in X and hence, we may assume $\hat{u}_n \rightarrow \hat{u}$, weakly in $H^1_0(\Omega)$ and strongly in $L^2(\Omega)$ to some $\hat{u} \in X$ such that $\|\hat{u}\|_{L^2} = 1$.

Choosing $\phi = u_n$ in (2.2), combining with (2.1) and using (2.3) we obtain

$$(2.4) \quad \left(\frac{1}{2} - \frac{1}{p+1}\right) \int_\Omega h|u_n|^{p+1} = o(1) \|u_n\|_{L^2}.$$

It follows from (2.4) and Fatou’s Lemma that

$$(2.5) \quad \int_\Omega h|\hat{u}|^{p+1} = 0$$

and hence $\hat{u} = 0$ a.e. on $\Omega \setminus \Omega_0$, that is, $\hat{u} \in H^1_*(\Omega_0)$. Fix $\phi \in C^\infty_0(\Omega)$, and observe that

$$(2.6) \quad \left| \int_\Omega h|u_n|^{p-1} u_n \phi \right| \leq c \|\phi\|_{L^\infty} \left(\int_\Omega h|u_n|^{p+1} \right)^{p/(p+1)}.$$

It follows from (2.2), (2.5) and (2.6) that

$$\int_{\Omega} \nabla \hat{u} \nabla \phi - \lambda \int_{\Omega} \hat{u} \phi = 0.$$

Hence \hat{u} is regular and satisfies

$$(2.7) \quad \Delta \hat{u} + \lambda \hat{u} = 0 \quad \text{in } \Omega.$$

But $\hat{u} \equiv 0$ on $\Omega \setminus \Omega_0$; thus the unique continuation property of the solutions of (2.7) implies $\hat{u} \equiv 0$ on Ω . This is a contradiction since $\|\hat{u}\|_{L^2} = 1$; then $\{u_n\}$ is bounded in $H_0^1(\Omega)$. Thus we can assume that the sequence $\{u_n\}$ is weakly convergent in $H_0^1(\Omega)$, strongly in $L^2(\Omega)$, and also a.e., to some $u \in H_0^1(\Omega)$. In order to show that the sequence converges in X it suffices to show that $\|u_n\|_X$ converges to $\|u\|_X$.

From (2.2) with $\phi = u_n$ we obtain

$$(2.8) \quad \lim_{n \rightarrow \infty} \int_{\Omega} |\nabla u_n|^2 + h|u_n|^{p+1} = \lambda \int_{\Omega} u^2.$$

Next, writing $J_\lambda = I + K$ where

$$I(u) = \frac{1}{2} \int_{\Omega} |\nabla u|^2 + \frac{1}{p+1} \int_{\Omega} h|u|^{p+1}$$

and using the convexity of I we see that for every $v \in X$

$$(I'(u_n) - I'(v))(u_n - v) = (-K'(u_n) + o(1) - I'(v))(u_n - v) \geq 0.$$

Choosing v appropriately and letting $n \rightarrow \infty$ we conclude that

$$(2.9) \quad \int_{\Omega} |\nabla u|^2 + \int_{\Omega} h|u|^{p+1} = \lambda \int_{\Omega} u^2.$$

On the other hand, we have

$$\int_{\Omega} |\nabla u|^2 \leq \liminf_{n \rightarrow \infty} \int_{\Omega} |\nabla u_n|^2$$

and

$$\int_{\Omega} h|u|^{p+1} \leq \liminf_{n \rightarrow \infty} \int_{\Omega} h|u_n|^{p+1}.$$

From these inequalities and (2.8), (2.9), we finally obtain that $\|u_n\|_X$ converges to $\|u\|_X$, as desired. \square

3. PROOF OF THEOREM 0.2

In this section we will prove that the functional J_λ defined in (0.5) possesses at least $s - k$ pairs of critical points when we are under the assumption

$$(3.1) \quad \lambda_s(\Omega) < \lambda < \lambda_{k+1}(\Omega_0)$$

with $s \geq k + 1$.

For this purpose we will use a variant of a result of Clark [2]; see also Rabinowitz [7]. Actually the result we need is a version of a theorem of Benci, Theorem 0.1 in [1]. For the reader's convenience we recall the notion of the genus, enumerate its main properties and sketch the proof of the critical point theorem.

We consider the Banach space X . We denote by \mathcal{E} the class of closed subsets A of $X \setminus \{0\}$ that are symmetric in the sense that $x \in A$ implies $-x \in A$. If $A \in \mathcal{E}$, then the genus of A is k if there is a continuous odd function

$$\varphi: A \rightarrow \mathbb{R}^k \setminus \{0\}$$

and k is the smallest integer with this property. If such a k does not exist, we say that the genus of A is ∞ . We write $\gamma(A)$ for the genus of A . In what follows we denote by $N_\delta(A)$ the uniform δ -neighborhood of the subset A of X , i.e. $N_\delta = \{x \in X \mid \|x - A\| \leq \delta\}$.

Proposition 3.1. *If $A, B \in \mathcal{E}$, then:*

- (1) *Normalization: If $x \neq 0$, then $\gamma(\{x\} \cup \{-x\}) = 1$.*
- (2) *Monotonicity: If $\varphi: A \rightarrow B$ is a continuous odd function, then $\gamma(A) \leq \gamma(B)$. In particular if $A \subset B$, then $\gamma(A) \leq \gamma(B)$.*
- (3) *Subadditivity: $\gamma(A \cup B) \leq \gamma(A) + \gamma(B)$.*
- (4) *Continuity: If A is compact, $\gamma(A) < \infty$ and there exists $\delta > 0$ so that $N_\delta(A) \in \mathcal{E}$ and $\gamma(A) = \gamma(N_\delta(A))$.*
- (5) *Dimension: If $A \in \mathcal{E}$, \mathcal{V} is a bounded neighborhood of 0 in \mathbb{R}^k , and there exists an odd homeomorphism between A and $\partial\mathcal{V}$, then $\gamma(A) = k$.*

For a proof of these properties we refer the reader to [7, Chapter 7]. Using these properties the following lemma can be proved (see Lemma 2.6 in [1]).

Lemma 3.1. *If $A \in \mathcal{E}$ is compact and Y is a subspace of X of finite codimension, then $\gamma(A \cap Y) \geq \gamma(A) - \text{codim}(Y)$.*

Proof. Let V be a vector subspace of X so that $X = V \oplus Y$ and consider $P: X \rightarrow V$ the projection onto V . Since A is compact, the continuity property implies that there exists $\delta > 0$ so that $\gamma(A \cap Y) = \gamma(N_\delta(A \cap Y))$. Put $N = N_\delta(A \cap Y)$.

Denote by s the codimension of Y and assume that

$$\gamma(A \cap Y) < \gamma(A) - s.$$

If we put $A_1 = A \cap N$ and $A_2 = \overline{A \setminus N}$, then $0 \notin P(A_2)$ and then, by the monotonicity and dimension properties of the genus,

$$\gamma(A_2) \leq \gamma(P(A_2)) \leq s.$$

On the other hand, using monotonicity and subadditivity we obtain

$$\gamma(A) \leq \gamma(A_1) + \gamma(A_2) \leq \gamma(N) + \gamma(A_2) < \gamma(A) - s + s,$$

a contradiction from which the result follows. \square

Next we formulate the critical point theorem to be used in our specific situation.

Theorem 3.1. *Let X be a real Banach space. Let $I: X \rightarrow \mathbb{R}$ be a C^1 functional satisfying the Palais-Smale condition. Assume further that*

- (1) *I is even and $I(0) = 0$.*
- (2) *There is a closed subspace Y of X so that I is bounded below on Y .*
- (3) *There is a closed subset $S \in \mathcal{E}$ and a constant $b > 0$ so that $I(w) \leq -b$, $\forall w \in S$, and*

(4) If $\mathcal{S}^h = h(S) \cap Y$, then $\gamma(\mathcal{S}^h) \geq m$ for all functions h in the class of functions

$$\Gamma = \{h: S \rightarrow X \setminus \{0\} | h \text{ is continuous and odd}\}.$$

Then the functional I possesses at least m pairs of critical points and their critical values have a minimax characterization (see (3.2)).

Proof. Since the proof is rather standard, we will be sketchy. First we define a class of sets. Given $1 \leq l \leq m$ we consider the classes \mathcal{A}_l defined as follows: $A \in \mathcal{A}_l$ if and only if $A = h(S) \setminus K$ such that $h \in \Gamma$, $\gamma(K) \leq m - l$ and $K \in \mathcal{E}$ is compact. By the monotonicity property of genus we have that $\mathcal{A}_m \subset \mathcal{A}_{m-1} \subset \dots \subset \mathcal{A}_1$.

For $1 \leq l \leq m$ we define the numbers

$$(3.2) \quad c_l = \inf_{A \in \mathcal{A}_l} \sup_{u \in A} I(u).$$

Since the classes \mathcal{A}_l are ordered, we have $c_1 \leq c_2 \leq \dots \leq c_m$. First we see that $c_1 > -\infty$. From the intersection assumption (4) and the subadditivity property we have that, for $1 \leq l \leq m$, $A \in \mathcal{A}_l$ implies $A = h(S) \setminus K$ and $\gamma(\overline{A \cap Y}) = \gamma(\overline{\mathcal{S}^h \setminus K}) \geq l$. This and assumption (2) yield $c_1 > -\infty$. It is also clear that $c_m < 0$, since we can take $S \in \mathcal{A}_m$. In order to show that the numbers c_l are critical values we proceed from here in the standard way. Only note that the Deformation Lemma will provide an odd homeomorphism $\eta: E \rightarrow E$ and if $h \in \Gamma$, then $\eta \circ h$ also belongs to Γ . See [7]. \square

In what follows we will prove that in our situation we can use Theorem 3.1. For a small number $\varepsilon > 0$, to be fixed later, we choose $\psi_1^\varepsilon, \dots, \psi_k^\varepsilon$ as in Lemma 1.1, part (a) and set

$$V_1 = \text{span}\{\psi_1^\varepsilon, \dots, \psi_k^\varepsilon\} \quad \text{and} \quad V_2 = \text{span}\{\psi_{k+1}, \dots, \psi_s\}.$$

Also we define

$$Y = \left\{ y \in X \mid \int_{\Omega} \nabla y \nabla \phi_i = 0, i = 1, \dots, k \right\}.$$

Then, by construction, we have that $V_1 \cap Y = \{0\}$, and hence $X = V_1 \oplus Y$. For small $\varepsilon > 0$ we also have $V_1 \cap V_2 = \{0\}$.

Lemma 3.2. *The functional J_λ defined in (0.5) satisfies*

- (1) J_λ is bounded below in Y , and
- (2) There exist numbers $b, \rho > 0$ such that

$$J_\lambda(w) \leq -b \quad \forall w \in V_1 \oplus V_2 \text{ with } \|w\|_X = \rho.$$

Proof. In order to prove (1) it suffices to verify that

$$J_\lambda(u) \rightarrow +\infty \quad \text{if } \|u\|_{H_0^1} \rightarrow +\infty, u \in Y.$$

Assume the contrary, that is, the existence of $c > 0$ and a sequence $\{u_n\} \subset Y$ with $\|u_n\|_{H_0^1} \rightarrow +\infty$ so that $J_\lambda(u_n) \leq c$, namely

$$(3.3) \quad \frac{1}{2} \int_{\Omega} |\nabla u_n|^2 - \lambda u_n^2 + \frac{1}{p+1} \int_{\Omega} h |u_n|^{p+1} \leq c.$$

We observe that $\|u_n\|_{L^2} \rightarrow +\infty$ and if we let $\hat{u}_n = u_n / \|u_n\|_{L^2}$, then

$$(3.4) \quad \frac{1}{2} \int_{\Omega} (|\nabla \hat{u}_n|^2 - \lambda) + \|u_n\|_{L^2}^{p-1} \left(\frac{1}{p+1} \int_{\Omega} h |\hat{u}_n|^{p+1} \right) \leq \frac{c}{\|u_n\|_{L^2}^2}.$$

Hence \hat{u}_n is bounded in $H_0^1(\Omega)$ and we may assume that $\hat{u}_n \rightharpoonup \hat{u}$ weakly in $H_0^1(\Omega)$ and strongly in $L^2(\Omega)$. From (3.4), using Fatou's Lemma we obtain $\int_{\Omega} h|\hat{u}|^{p+1} = 0$; hence $\hat{u} \in H_*^1(\Omega_0)$. But \hat{u} also belongs to Y , the H_0^1 -orthogonal of $\text{span}\{\phi_1, \dots, \phi_k\}$, hence to the H_*^1 -orthogonal of this space. On the other hand (3.4) also yields

$$\int_{\Omega} |\nabla \hat{u}|^2 \leq \lambda$$

which, by the variational characterization of the eigenvalues, implies $\lambda \geq \lambda_{k+1}(\Omega_0)$. This contradicts our hypothesis (3.1) and (1) is therefore proved.

Next we prove (2). Let

$$w = \sum_{i=1}^s t_i(\psi_i + \eta_i^\varepsilon)$$

where $\eta_i^\varepsilon = \psi_i^\varepsilon - \psi_i$ if $i = 1, \dots, k$ and $\eta_i^\varepsilon = 0$ if $i = k + 1, \dots, s$. We compute

$$(3.5) \quad J_\lambda(w) = \frac{1}{2} \sum_{i=1}^s t_i^2(\lambda_i(\Omega) - \lambda) + \frac{1}{p+1} \int_{\Omega} h|w|^{p+1}$$

$$(3.6) \quad + \frac{1}{2} \sum_{i=1}^s \sum_{j=1}^s t_i t_j \left\{ \int_{\Omega} \nabla(\psi_i + \eta_i^\varepsilon) \nabla \eta_j^\varepsilon - \lambda(\psi_i + \eta_i^\varepsilon) \eta_j^\varepsilon \right\}.$$

Hence, letting $|w|^2 = \sum_{i=1}^s t_i^2$ we obtain

$$(3.7) \quad J_\lambda(w) \leq -c_1|w|^2 + c_2\varepsilon|w|^2 + c_3|w|^{p+1}$$

for certain positive constants c_1, c_2 and c_3 , independent of ε . From (3.7), the result of (2) readily follows after choosing ε sufficiently small. \square

In the rest of this section we will denote

$$S_\rho = \{w \in V_1 \oplus V_2 \mid \|w\|_X = \rho\}$$

where ρ is chosen as in Lemma 3.2. First we define the class of functions

$$\Gamma = \{h: S_\rho \rightarrow X \setminus \{0\} \mid h \text{ is continuous and odd}\}.$$

We note that if $\eta: X \rightarrow X$ is an odd homeomorphism, then $\eta \circ h \in \Gamma$ for all $h \in \Gamma$.

The next lemma gives an intersection result that implies hypothesis (4) in Theorem 3.1.

Lemma 3.3. *For $h \in \Gamma$ we let $\mathcal{S}^h = h(S_\rho) \cap Y$. Then $\gamma(\mathcal{S}^h) \geq s - k$.*

Proof. Since $S_\rho \subset V_1 \oplus V_2$ is finite dimensional, $A = h(S_\rho)$ is compact. By the dimension property of the genus, $\gamma(A) = s$. Then Lemma 3.1 gives the result. \square

Proof of Theorem 0.2. Lemmas 3.2 and 3.3 imply that Theorem 3.1 is applicable to the functional J_λ when λ satisfies (0.4). Thus there exist pairs of solutions $\pm u_\lambda^{k+l}$, $l = 1, \dots, s - k$, such that

$$J_\lambda(u_\lambda^{k+l}) = c_l(\lambda) = \inf_{A \in \mathcal{A}_l} \sup_{u \in A} J_\lambda(u), \quad l = 1, \dots, s - k,$$

where the class of sets \mathcal{A}_l was defined in the proof of Theorem 3.1. Let q be the multiplicity of $\lambda_k(\Omega_0)$ and denote $\bar{q} = \min\{q, s - k\}$. It is easily checked that the validity of assertions (i) and (ii) of Theorem 0.2 will follow if we prove the following two facts:

Claim 1. $c_{\bar{q}}(\lambda) \rightarrow -\infty$ as $\lambda \rightarrow \lambda_{k+1}(\Omega_0)$.

Claim 2. If l satisfies $q < l \leq s - k$, then there is a real number c such that $c_l(\lambda) \geq c$ for all λ satisfying (0.4).

For the proof of Claim 1, we need some estimates for J_λ which we carry out next. In the rest of the proof we will assume $\bar{q} = q < s - k$, the other case being similar.

Note that the functions $\phi_{k+1}, \dots, \phi_{k+q}$ are not solutions of the equation

$$(3.8) \quad -\Delta u = \lambda_{k+1}(\Omega_0)u \quad \text{in } \Omega,$$

$$(3.9) \quad u = 0 \quad \text{on } \partial\Omega;$$

hence the functionals $f_i \in H^{-1}(\Omega)$, $i = 1, \dots, q$, defined as

$$(3.10) \quad f_i(\varphi) = \int_{\Omega} \nabla \phi_{k+i} \nabla \varphi - \lambda_{k+1}(\Omega_0) \phi_{k+i} \varphi \quad \forall \varphi \in H_0^1(\Omega)$$

are linearly independent. This implies the existence of functions $v_1, \dots, v_q \in H_0^1(\Omega)$ so that the matrix $A = \{a_{ij}\}$ defined as

$$(3.11) \quad a_{ij} = f_i(v_j), \quad 1 \leq i, j \leq q,$$

is invertible. Moreover, we may assume without loss of generality that the functions v_j satisfy

$$(3.12) \quad \int_{\Omega} \nabla \phi_l \nabla v_j = 0, \quad 1 \leq j \leq q$$

and $1 \leq l \leq k$, $k + q + 1 \leq l \leq s$.

Let us consider $w_1 \in \text{span}\{\phi_1, \dots, \phi_k\}$ and $w_2 \in \text{span}\{\phi_{k+q+1}, \dots, \phi_s\}$. For $\theta, t \in \mathbb{R}^q$ with components θ_i and t_i , $1 \leq i \leq q$, define $v = \sum_{i=1}^q \theta_i v_i$ and $u = \sum_{i=1}^q t_i \phi_{i+k}$. We want to estimate $J_\lambda(w_1 + v + u + w_2)$. Using the definition of J_λ we find that

$$(3.13) \quad \mathcal{J} \equiv J_\lambda(w_1 + v + u + w_2) = \mathcal{J}_1 + \mathcal{J}_2,$$

where

$$(3.14) \quad \mathcal{J}_1 = J_\lambda(w_1) + J_\lambda(w_2) + J_\lambda(v) - \lambda \int_{\Omega} (w_1 + w_2)v,$$

and

$$(3.15) \quad \mathcal{J}_2 = \frac{\lambda_{k+1}(\Omega_0) - \lambda}{2} |t|^2 + \sum_{i=1}^q \sum_{j=1}^q \theta_i t_j \int_{\Omega} \nabla v_i \nabla \phi_{j+k} - \lambda v_i \phi_{j+k}.$$

Using the matrix A defined in (3.11) we rewrite this last term as

$$(3.16) \quad \mathcal{J}_2 = \frac{\lambda_{k+1}(\Omega_0) - \lambda}{2} |t|^2 + (\lambda_{k+1}(\Omega_0) - \lambda) \sum_{i=1}^q \sum_{j=1}^q \theta_i t_j \int_{\Omega} v_i \phi_{j+k} + (\theta, A^T t)$$

where $|\cdot|$ and (\cdot, \cdot) denote the usual norm and inner product in \mathbb{R}^q , and A^t represents the transpose of A . Then, assuming that θ is bounded, from (3.16) we obtain

$$(3.17) \quad \mathcal{J}_2 \leq \frac{\lambda_{k+1}(\Omega_0) - \lambda}{2} (|t|^2 + a_1|t|) + (\theta, A^t t).$$

Here and henceforth we use the notation a_1, a_2 , etc., to designate different positive absolute constants. On the other hand, assuming that θ is bounded, from (3.14) we have that

$$(3.18) \quad \mathcal{J}_1 \leq \frac{\lambda_k(\Omega_0) - \lambda}{2} \|w_1\|_{L^2}^2 + a_1 \|w_2\|_{L^2}^2 + a_2 \|w_1 + w_2\|_{L^2} + a_3.$$

Next we will find an appropriate function h in Γ and a set K with $\gamma(K) \geq (s - k) - q$ in order to estimate the critical value $c_q(\lambda)$ when λ is close to $\lambda_{k+1}(\Omega_0)$. Let us fix a number $\sigma \geq 1$. We define the function $h_\sigma: V_1 \oplus V_2 \rightarrow X$ in the following manner. For $\alpha_i \in \mathbb{R}$, $1 \leq i \leq s$, we put

$$v_1 = \sum_{i=1}^k \alpha_i \psi_i^\varepsilon, \quad \bar{v} = \sum_{i=k+1}^{k+q} \alpha_i \psi_i, \quad v_2 = \sum_{i=k+q+1}^s \alpha_i \psi_i$$

and define

$$(3.19) \quad h_\sigma(v_1 + \bar{v} + v_2) = \sigma^{1/2} w_1 + v + \sigma u + w_2$$

where

$$w_1 = \sum_{i=1}^k \alpha_i \phi_i, \quad v = \sum_{i=k+1}^{k+q} \beta_i v_{i-k}, \quad u = \sum_{i=k+1}^{k+q} \alpha_i \phi_i, \quad w_2 = \sum_{i=k+q+1}^s \alpha_i \phi_i,$$

and the coefficients β_i are given by

$$(3.20) \quad \beta_i = - \sum_{j=1}^q a_{ji} \alpha_j.$$

Next define, for a small $\nu > 0$ and h_σ as in (3.19), the compact set

$$(3.21) \quad K_\nu = \{h_\sigma(v_1 + \bar{v} + v_2) \mid \|v_1 + \bar{v} + v_2\|_X = \rho, \|v_2\|_X \geq \nu\}.$$

It is easy to see that $\gamma(K_\nu) = \gamma(K_1)$ and then $\gamma(K_\nu) = s - (k + q) = (s - k) - q$. On the other hand, for $z \in h_\sigma(S_\rho) \setminus K_\nu$, equations (3.17), (3.18) and (3.20) imply that

$$(3.22) \quad \begin{aligned} J_\lambda(z) &\leq (\lambda_{k+1}(\Omega_0) - \lambda) a_1 \sigma^2 - \sigma a_2 \|\bar{v}\|_X^2 \\ &\quad + (\lambda_k(\Omega_0) - \lambda) \sigma \|v_1\|_{L^2}^2 + \sigma^{1/2} a_3 + a_4. \end{aligned}$$

If $\|\bar{v}\|_X \geq \rho/4$, from (3.22) we find that

$$(3.23) \quad J_\lambda(z) \leq (\lambda_{k+1}(\Omega_0) - \lambda) a_1 \sigma^2 - \sigma \rho a_2 + a_3 \sigma^{1/2} + a_4$$

and if $\|\bar{v}\|_X < \rho/4$, $\lambda \geq \tilde{\lambda} > \lambda_k(\Omega_0)$ and $\nu < \rho/4$, we have from (3.22) that

$$(3.24) \quad J_\lambda(z) \leq (\lambda_{k+1}(\Omega_0) - \lambda) a_1 \sigma^2 - a_2 \sigma \rho^2 + a_3 \sigma^{1/2} + a_4.$$

Now, combining (3.23) and (3.24) we observe that an appropriate choice of σ will yield

$$J_\lambda(z) \leq - \frac{b_1}{\lambda_{k+1}(\Omega_0) - \lambda} + b_2 \quad \forall z \in h_\sigma(S_\rho) \setminus K_\nu$$

for λ close to $\lambda_{k+1}(\Omega_0)$, where the positive constants b_1 and b_2 may depend on ρ but not on λ . The definition of $c_q(\lambda)$ as a minimax implies that

$$(3.25) \quad c_q(\lambda) \leq \sup\{J_\lambda(z) \mid z \in h_\sigma(S_\rho) \setminus K_\nu\}$$

from which the validity of Claim 1 follows at once.

Next we prove Claim 2. If we define

$$Y_q = \left\{ y \in X \mid \int_\Omega \nabla y \nabla \phi_i = 0, 1 \leq i \leq k+q \right\},$$

then the same argument given for the proof of Lemma 3.3 will imply that

$$(3.26) \quad \gamma(\mathcal{S}^{h,q}) \geq s - (k+q) \quad \forall h \in \Gamma$$

where $\mathcal{S}^{h,q} = h(S_\rho) \cap Y_q$. Consequently, for $l \geq q+1$ we will have that

$$\gamma(\overline{h(S_\rho) \setminus K \cap Y_q}) \geq l - q \geq 1.$$

Thus, for every $A \in \mathcal{A}_l$ there is $y \in A$ such that $y \in Y_q$. Following the estimates that led us to Lemma 3.2, part (1) we will find that J_λ is bounded below by a constant independent of λ , for λ satisfying (0.4). Thus $c_l(\lambda) \geq c$ for all $q+1 \leq l \leq s$ if $\lambda \leq \lambda_{k+1}(\Omega_0)$, as desired. Thus Claim 2 is established, and the theorem follows. \square

Remark 3.1. It should be noticed that the above proof does not require homogeneity of the nonlinearity, and will therefore apply to the more general problem (1.14)–(1.15) as long as f is odd, increasing and satisfies appropriate polynomial growth restrictions near 0 and ∞ .

REFERENCES

1. V. Benci, *On critical point theory for indefinite functionals in the presence of symmetries*, Trans. Amer. Math. Soc. **274** (1982), 533–572.
2. D. Clark, *A variant of the Lusternik-Schnirelman theory*, Indiana Univ. Math. J. **22** (1972), 65–74.
3. M. del Pino, *Positive solutions of a semilinear elliptic equation on a compact manifold* (to appear).
4. J. Kazdan and F. Warner, *Scalar curvature and conformal deformation of Riemannian structure*, J. Differential Geom. **10** (1975), 113–134.
5. T. Ouyang, *On the positive solutions of semilinear equations $\Delta u + \lambda u - hu^p = 0$ on compact manifolds*, Trans. Amer. Math. Soc. **331** (1992), 503–527.
6. ———, *On the positive solutions of semilinear equations $\Delta u + \lambda u - hu^p = 0$ on compact manifolds*, Part II, Indiana Univ. Math. J. **40** (1991), 1083–1140.
7. P. Rabinowitz, *Minimax methods in critical point theory with applications to differential equations*, C.B.M.S. Regional Conf. Ser. in Math., no. 65, American Mathematical Society, Providence, RI, 1986.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, CHICAGO, ILLINOIS 60637
 E-mail address: delpino@math.uchicago.edu

DEPARTAMENTO DE INGENIERIA MATEMÁTICA, F.C.F.M., UNIVERSIDAD DE CHILE, CASILLA 170
 CORREO 3, SANTIAGO, CHILE
 E-mail address: pfelmer@llaima.dim.uchile.cl